




NORTHWESTERN
UNIVERSITY



THE DEVELOPMENT OF A HUMANIZED ANTIBODY-TARGETED ABO-SPECIFIC PET PROBE FOR EARLY DIAGNOSTIC IMAGING OF ALZHEIMER'S DISEASE

Adrian Bebenek

Mentors: Ms. Kirsten Viola and Dr. William Klein

Kirsten L. Viola¹, Adrian Bebenek¹, Jake Vitrofsky¹, Maira Bicca², Erika N. Cline¹, Alex L. Qin¹, Chad Haney¹, Ting-Tung Chang³, and William Klein^{1, 4}

¹ Northwestern University

² Universidade Federal de Santa Catarina

³ Rethink Imaging

⁴ Acumen Pharmaceuticals

TOPICS

- Topic 1 – General Background
- Topic 2 – Molecular Basis
- Topic 3 – Design of PET Probe
- Topic 4 – Preservation of Immunoreactivity
- Topic 5 – NU4PET Preliminary Injections
- Topic 6 – NU4 Histology and Morphology
- Topic 7 – ACU193 Histology and Morphology
- Topic 8 – ACU193PET Preliminary Injections

BACKGROUND



It's the only cause of death
in the top 10 in America that
**CANNOT BE PREVENTED,
CURED OR SLOWED.**



ALMOST TWO THIRDS
of Americans with Alzheimer's
disease are women.

1 IN 3

SENIORS
dies with Alzheimer's or
another dementia.

6
Alzheimer's
disease is the
**6TH LEADING
CAUSE OF DEATH
IN THE UNITED
STATES.**

Only

45%

of people with
ALZHEIMER'S
disease or their
caregivers report
**BEING TOLD OF
THEIR DIAGNOSIS.**



More than

90%

of people with the
four most common
types of **CANCER**
have been
**TOLD OF THEIR
DIAGNOSIS.**



By 2050, these costs
could rise as high as
\$1.1 TRILLION.




In 2015, Alzheimer's and
other dementias will cost the
nation **\$226 BILLION.**


Alzheimers
Association 2015

MOLECULAR BASIS

- Amyloid- β plaques
- Neurofibrillary tangles




PBS 2016



Lieff 2015


AMYLOID BETA PLAQUES

- Amyloid Precursor Protein (APP)
- $A\beta_{42}$ vs $A\beta_{40}$
- Peptide \rightarrow oligomer \rightarrow fibril \rightarrow plaque




AMYLOID CASCADE HYPOTHESIS

- Weak correlation between AD and plaques
- Amyloid- β oligomers ($A\beta$ Os) initiate neurodegeneration




NIA 2016

BACKGROUND



Medical Care
Corporation
2015

ANTIBODIES




Cell Applications 2015

- Monoclonal
- NU4 - mouse
- ACU193 - human


PET SYNTHESIS

- PET uses radioactive ligand
- Conjugation of DOTA cage to antibody
- Chelation with radioactive compound/element (^{64}Cu)



ABO ARE SEPARATE FROM PLAQUES


- Demonstrate that A β O are separate from plaques
- Immunostained tg and wt mice with 568-NU4 (A β O)
- Counterstained with ThioFlavin S (amyloid plaques)



Maira Bicca



DOTA BINDS TO THE FC REGION OF NU4

- Electrophoresis
- Confirm DOTA binds to constant region
- NU4: 54.091 kDa
- NU4-DOTA: 56.849 kDa
- 6-7 DOTA bound to heavy chain




NU4DOTA RETAINS IMMUNOREACTIVITY TO ABOS

- Dot blots
- EC_{50}
- No decrease in immunoreactivity




NU4-DOTA COLOCALIZES WITH ABOS *IN VITRO*

- NU4DOTA and FAM-A β Os in primary hippocampal neuron culture
- High levels of colocalization
- Retained immunoreactivity *in vitro*



60x


FAM ADDLs/NU4



Maira Bicca

NU4PET DEMONSTRATE A STRONG AD DEPENDENT SIGNAL IN MICE


- Tg and Wt mice injected
- NU4PET
- IgGPET
- NU4PET demonstrates a strong AD dependent signal



Ting-Tung Chang

NU4PET DEMONSTRATES SUBSTANTIAL BRAIN UPTAKE


- Blood brain barrier is extremely difficult to cross
- Percent of injected dose retained similar to PiB and Florbetapir



Ting-Tung Chang


PET SIGNAL CORRELATES TO ACU193 IMMUNOFLUORESCENT INTENSITY

- Immunofluorescently labeled brains with ACU193
- Fluorescent intensity correlates to PET signal




ABOS ARE DETECTABLE AT 2 MONTHS IN 5XFAD MICE

- Target for diagnosis must appear before onset of symptoms
- ERENNA and histology
- A β O_s are detectable at 2 months




Erika Cline




NU4 DEMONSTRATES PLAQUE-LIKE AND PUNCTATE LABELING OF PYRAMIDAL LAYER IN 5XFAD MOUSE MODEL



- Identify pathology seen in our 5xFAD mouse model
- Punctate and plaque-like labeling of the dendritic arbors
- A β O_s found primarily in hippocampus, dentate gyrus and along the pyramidal layer







ACU193 DISTINGUISHES BETWEEN DISEASED AND NONDISEASED HUMAN BRAINS




ACU193 SHOWS PLAQUE-LIKE, PUNCTATE, AND DIFFUSE PATHOLOGY IN HUMAN BRAIN TISSUE



ACU193DOTA DISTINGUISHES BETWEEN 5XFAD AND WT TISSUES DOWN TO 2 MONTHS (NU4 AND ACU193DOTA)




ACU193DOTA BINDS IN AN AGE DEPENDENT MANNER AND DISTINGUISHES BETWEEN TG AND WT MICE




ACU193PET SHOWS PROMISE IN DISTINGUISHING BETWEEN DISEASED AND NONDISEASED MICE

DAY 1 RESULTS

Wild Type




5xFAD




DAY 1 RESULTS

Wild Type



5xFAD



CONCLUSION

- ACU193PET demonstrates tremendous potential as an early diagnostic imaging tool for AD
- Future Work:
 - Substantiating AD specific probe signal by autoradiography using mouse and human brain sections
 - Dose curve PET to determine optimal dosing range in mid-stage AD (5xFAD model)
 - Evaluation of gross uptake and clearance in mice in vivo
 - Longitudinal analysis to determine earliest stages at which PET probe detects A β Os
 - Quantitative relationship between PET signals, A β Os detected histologically, A β Os detected biochemically, and memory loss
 - Human trial in 18 months
 - Dual MRI/PET

ACKNOWLEDGEMENTS

First and foremost I would like to thank Ms. Kirsten Viola for allowing me to work on this project with her and for being my mentor. I would like to thank Dr. William Klein for the opportunity to work in his lab and for all of the mentorship that he has also provided. I would also like to thank everyone who I have worked with at Klein Lab and all collaborators outside of Northwestern University for making this possible. Finally, would like to thank the SIR staff at IMSA and my parents for providing me with this opportunity and being a tremendous source of support.

- NIH NIA Award Number AG045637
- NCI CCSG P30 CA060553 and NIH 1S10OD010398-01 for Northwestern University's Center for Advanced Microscopy
- NCI CCSG P30 CA060553 for Northwestern University's Center for Advanced Molecular Imaging

BIBLIOGRAPHY

1. Alzheimer's Association. (2013, September 16). Latest Alzheimer's Facts and Figures. Retrieved June 19, 2015, from <http://www.alz.org/facts/overview.asp>
2. Slemmon, J. R. et al. Measurement of Aβ₁₋₄₂ in cerebrospinal fluid is influenced by matrix effects. *Journal of neurochemistry* 120, 325-333, doi:10.1111/j.1471-4159.2011.07553.x (2012).
3. Landau, S. M., et al. (2013). Amyloid-beta imaging with Pittsburgh compound B and florbetapir: comparing radiotracers and quantification methods. *J Nucl Med.* 54(1): 70-77. doi: 10.2967/jnumed.112.109009.
4. Braak, H. & Braak, E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol.* 82, 239–259 (1991). Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/1759558>
5. Tomiyama, T. et al. (2010). A mouse model of amyloid beta oligomers: their contribution to synaptic alteration, abnormal tau phosphorylation, glial activation, and neuronal loss in vivo. *J. Neurosci.* 30, 4845–4856. doi: 10.1523/JNEUROSCI.5825-09.2010.
6. Tomic, J.L., Pensalfini, A., Head, E. & Glabe, C.G. (2009). Soluble fibrillar oligomer levels are elevated in Alzheimer's disease brain and correlate with cognitive dysfunction. *Neurobiol. Dis.* 35, 352–358. Retrieved from: doi: 10.1016/j.nbd.2009.05.024.
7. Viola, K.L., Sbarboro, J., Sureka, R., De, M., Bicca, M.A., ...& Klein, W.L. (2014). Towards non-invasive diagnostic imaging of early-stage Alzheimer's disease. *Nature Nanotechnology.* 10, 91-98. doi: 10.1038/nnano.2014.254
8. Lambert, M.P., Velasco, P.T., Chang, L., Viola, K.L., Fernandez, S., ...Klein, W.L. (2007). Monoclonal antibodies that target pathological assemblies of Aβ. *Journal of Neurochemistry.* 100(1), 23-35. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/17116235>